

Our Ref.: CL001066-CIP

U.S. PATENT APPLICATION

Inventor(s): Ming-Hui WEI
Karen A. KETCHUM
Valentina DI FRANCESCO
Ellen M. BEASLEY

Invention: ISOLATED HUMAN PHOSPHATASE PROTEINS, NUCLEIC ACID
MOLECULES ENCODING HUMAN PHOSPHATASE PROTEINS, AND
USES THEREOF

CELERA GENOMICS CORPORATION.
45 WEST GUDE DR., C2-4#20
ROCKVILLE, MD 20850
(240) 453-3067
Fax (240)-453-3084

SPECIFICATION

**ISOLATED HUMAN PHOSPHATASE PROTEINS, NUCLEIC ACID
MOLECULES ENCODING HUMAN PHOSPHATASE PROTEINS, AND USES
THEREOF**

5 RELATED APPLICATIONS

The present application is a Continuation-In-Part of US Serial No. 09/752,820, filed January 3, 2001 (Atty. Docket CL001066).

FIELD OF THE INVENTION

10 The present invention is in the field of phosphatase proteins that are related to the PPM1G phosphatase subfamily, recombinant DNA molecules and protein production. The present invention specifically provides novel phosphatase peptides and proteins and nucleic acid molecules encoding such peptide and protein molecules, all of which are useful in the development of human therapeutics and diagnostic compositions and methods.

BACKGROUND OF THE INVENTION

15 Phosphatase proteins, particularly members of the PPM1G phosphatase subfamily, are a major target for drug action and development. Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown members of this subfamily of phosphatase proteins. The present invention advances the state of the art by providing a previously unidentified human phosphatase proteins that have homology to members of the PPM1G phosphatase subfamily.

Protein Phosphatase

25 Cellular signal transduction is a fundamental mechanism whereby external stimuli that regulate diverse cellular processes are relayed to the interior of cells. The biochemical pathways through which signals are transmitted within cells comprise a circuitry of directly or functionally connected interactive proteins. One of the key biochemical mechanisms of signal transduction involves the reversible phosphorylation

of certain residues on proteins. The phosphorylation state of a protein may affect its conformation and/or enzymic activity as well as its cellular location. The phosphorylation state of a protein is modified through the reciprocal actions of protein kinases (PKs) and protein phosphatases (PPs) at various specific amino acid residues.

5 Protein phosphorylation is the ubiquitous strategy used to control the activities of eukaryotic cells. It is estimated that 10% of the proteins active in a typical mammalian cell are phosphorylated. The high-energy phosphate that confers activation and is transferred from adenosine triphosphate molecules to protein-by-protein phosphatases is subsequently removed from the protein-by-protein phosphatases. In this way, the
10 phosphatases control most cellular signaling events that regulate cell growth and differentiation, cell-to-cell contacts, the cell cycle, and oncogenesis.

The protein phosphorylation/dephosphorylation cycle is one of the major regulatory mechanisms employed by eukaryotic cells to control cellular activities. It is estimated that more than 10% of the active proteins in a typical mammalian cell are phosphorylated. During protein phosphorylation/dephosphorylation, phosphate groups are transferred from adenosine triphosphate molecules to protein-by-protein phosphatases and are removed from the protein-by-protein phosphatases.

Protein phosphatases function in cellular signaling events that regulate cell growth and differentiation, cell-to-cell contacts, the cell cycle, and oncogenesis. Three protein phosphatase families have been identified as evolutionarily distinct. These include the serine/threonine phosphatases, the protein tyrosine phosphatases, and the acid/alkaline phosphatases (Carbonneau H. and Tonks N. K. (1992) *Annu. Rev. Cell Biol.* 8:463-93).

The serine/threonine phosphatases are either cytosolic or associated with a receptor. On the basis of their sensitivity to two thermostable proteins, inhibitors 1 and 2,
25 and their divalent cation requirements, the serine/threonine phosphatases can be separated into four distinct groups, PP-I, PP-IIA, PP-IIB, and PP-IIC.

PP-I dephosphorylates many of the proteins phosphorylated by cyclic AMP-dependent protein kinase and is therefore an important regulator of many cyclic AMP mediated, hormone responses in cells. PP-IIA has broad specificity for control of
30 cell cycle, growth and proliferation, and DNA replication and is the main phosphatase responsible for reversing the phosphorylations of serine/threonine phosphatases. PP-IIB,

or calcineurin (Cn), is a Ca^{2+} -activated phosphatase; it is involved in the regulation of such diverse cellular functions as ion channel regulation, neuronal transmission, gene transcription, muscle glycogen metabolism, and lymphocyte activation.

5 PP-IIC is a Mg^{++} -dependent phosphatase which participates in a wide variety of functions including regulating cyclic AMP-activated protein-phosphatase activity, Ca^{++} -dependent signal transduction, tRNA splicing, and signal transmission related to heat shock responses. PP-IIC is a monomeric protein with a molecular mass of about 40-45 kDa. One α and several β isoforms of PP-IIC have been identified (Wenk, J. et al. (1992) FEBS Lett. 297: 135-138; Terasawa, T. et al. 10 (1993) Arch. Biochem. Biophys. 307: 342-349; and Kato, S. et al. (1995) Arch. Biochem. Biophys. 318: 387-393).

15 The levels of protein phosphorylation required for normal cell growth and differentiation at any time are achieved through the coordinated action of PKs and PPS. Depending on the cellular context, these two types of enzymes may either antagonize or cooperate with each other during signal transduction. An imbalance between these enzymes may impair normal cell functions leading to metabolic disorders and cellular transformation.

20 For example, insulin binding to the insulin receptor, which is a PTK, triggers a variety of metabolic and growth promoting effects such as glucose transport, biosynthesis of glycogen and fats, DNA synthesis, cell division and differentiation. Diabetes mellitus, which is characterized by insufficient or a lack of insulin signal transduction, can be caused by any abnormality at any step along the insulin signaling pathway. (Olefsky, 1988, in "Cecil Textbook of Medicine," 18th Ed., 2:1360-81).

25 It is also well known, for example, that the overexpression of PTKs, such as HER2, can play a decisive role in the development of cancer (Slamon et al., 1987, Science 235:77-82) and that antibodies capable of blocking the activity of this enzyme can abrogate tumor growth (Drebin et al., 1988, Oncogene 2:387-394). Blocking the signal transduction capability of tyrosine phosphatases such as Flk-1 and the PDGF receptor have been shown to block tumor growth in animal models (Millauer et al., 1994, 30 Nature 367:577; Ueno et al., Science, 252:844-848).

Relatively less is known with respect to the direct role of phosphatases in signal transduction; PPs may play a role in human diseases. For example, ectopic expression of RPTP.alpha. produces a transformed phenotype in embryonic fibroblasts (Zheng et al., Nature 359:336-339), and overexpression of RPTP.alpha. in embryonal carcinoma cells causes the cells to differentiate into a cell type with neuronal phenotype (den Hertog et al., EMBO J 12:3789-3798). The gene for human RPTP.gamma. has been localized to chromosome 3p21 which is a segment frequently altered in renal and small lung carcinoma. Mutations may occur in the extracellular segment of RPTP.gamma. which renders a RPTP that no longer respond to external signals (LaForgia et al., Wary et al., 1993, Cancer Res 52:478-482). Mutations in the gene encoding PTP1C (also known as HCP, SHP) are the cause of the moth-eaten phenotype in mice that suffer severe immunodeficiency, and systemic autoimmune disease accompanied by hyperproliferation of macrophages (Schultz et al., 1993, Cell 73:1445-1454). PTP1D (also known as Syp or PTP2C) has been shown to bind through SH2 domains to sites of phosphorylation in PDGFR, EGFR and insulin receptor substrate 1 (IRS-1). Reducing the activity of PTP1D by microinjection of anti-PTP1D antibody has been shown to block insulin or EGF-induced mitogenesis (Xiao et al., 1994, J Biol Chem 269:21244-21248).

Magnesium-Dependent Protein Phosphatase 1G (PPM1G)

The novel human protein, and encoding gene, provided by the present invention is a novel splice form of PPM1G, formerly referred to as PP2C-gamma, a member of the serine/threonine phosphatase family. Specifically, the novel PPM1G variant provided by the present invention is missing exon 5, exon 6, and exon 7 compared with the known PPM1G isoform (gi4505999).

PPM1G is magnesium or manganese dependent and is characterized by a large, unique acidic domain not found in other phosphatases (Travis *et al.*, *FEBS Lett.* 412 (3), 415-419 (1997)). PPM1G is a component of splicing-complementing factor-1 (SCF1) and plays an important role in pre-mRNA splicing and spliceosome assembly. It has been suggested that PPM1G catalyzes at least one specific dephosphorylation event critical for spliceosome assembly. This is further supported by the findings that PPM1G localizes to the nucleus, mutations in the active site required for phosphatase activity abolish

spliceosome activity, and substitution of PP2C-alpha, which is closely related to PPM1G (also referred to as PP2C-gamma), also abolishes splicing function (Murray *et al.*, *Genes Dev* 1999 Jan 1;13(1):87-97).

5 The discovery of a new human protein phosphatase and the polynucleotides encoding it satisfies a need in the art by providing new compositions that are useful in the diagnosis, prevention and treatment of biological processes associated with abnormal or unwanted protein phosphorylation.

10 **SUMMARY OF THE INVENTION**

15 The present invention is based in part on the identification of amino acid sequences of human phosphatase peptides and proteins that are related to the PPM1G phosphatase subfamily, as well as allelic variants and other mammalian orthologs thereof. These unique peptide sequences, and nucleic acid sequences that encode these peptides, can be used as models for the development of human therapeutic targets, aid in the identification of therapeutic proteins, and serve as targets for the development of human therapeutic agents that modulate phosphatase activity in cells and tissues that express the phosphatase. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye
20 retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes.

DESCRIPTION OF THE FIGURE SHEETS

25 FIGURE 1 provides the nucleotide sequence of a cDNA molecule or transcript sequence that encodes the phosphatase protein of the present invention. (SEQ ID NO:1) In addition, structure and functional information is provided, such as ATG start, stop and tissue distribution, where available, that allows one to readily determine specific uses of inventions based on this molecular sequence. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin

melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes.

FIGURE 2 provides the predicted amino acid sequence of the phosphatase of the present invention. (SEQ ID NO:2) In addition structure and functional information such as protein family, function, and modification sites is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence.

FIGURE 3 provides genomic sequences that span the gene encoding the phosphatase protein of the present invention. (SEQ ID NO:3) In addition, structure and functional information, such as intron/exon structure, promoter location, etc., is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence. As illustrated in Figure 3, 21 SNPs were identified. Figure 3 also provides an alignment, from the "Genewise" computer program, of the amino sequences of the present invention and gi4505999 (the top BLAST hit) against the genomic sequence of the present invention. This alignment further illustrates that the novel human phosphatase protein provided by the present invention is a novel splice variant that is missing exons 5, 6, and 7 compared to gi4505999.

DETAILED DESCRIPTION OF THE INVENTION

General Description

The present invention is based on the sequencing of the human genome. During the sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a phosphatase protein or part of a phosphatase protein and are related to the PPM1G phosphatase subfamily. Utilizing these sequences, additional genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of human phosphatase peptides and proteins that are related to the PPM1G phosphatase subfamily, nucleic acid sequences in the form of transcript sequences, cDNA sequences and/or genomic sequences that encode these phosphatase

peptides and proteins, nucleic acid variation (allelic information), tissue distribution of expression, and information about the closest art known protein/peptide/domain that has structural or sequence homology to the phosphatase of the present invention.

In addition to being previously unknown, the peptides that are provided in the present invention are selected based on their ability to be used for the development of commercially important products and services. Specifically, the present peptides are selected based on homology and/or structural relatedness to known phosphatase proteins of the PPM1G phosphatase subfamily and the expression pattern observed. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. The art has clearly established the commercial importance of members of this family of proteins and proteins that have expression patterns similar to that of the present gene. Some of the more specific features of the peptides of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for each of the known PPM1G family or subfamily of phosphatase proteins.

Specific Embodiments

Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being members of the phosphatase family of proteins and are related to the PPM1G phosphatase subfamily (protein sequences are provided in Figure 2, transcript/cDNA sequences are provided in Figure 1 and genomic sequences are provided in Figure 3). The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the phosphatase peptides of the present invention, phosphatase peptides, or peptides/proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprise the amino acid sequences of the phosphatase peptides disclosed in the Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The peptides of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components (the features of an isolated nucleic acid molecule is discussed below).

In some uses, "substantially free of cellular material" includes preparations of the peptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the phosphatase peptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

The isolated phosphatase peptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanomas, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung

small cell carcinomas, and leukocytes. For example, a nucleic acid molecule encoding the phosphatase peptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

Accordingly, the present invention provides proteins that consist of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid sequence when the amino acid sequence is the final amino acid sequence of the protein.

The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about 1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein comprises an amino acid sequence when the amino acid sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the phosphatase peptides of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

The phosphatase peptides of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a phosphatase peptide operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the phosphatase peptide. "Operatively linked" indicates that the phosphatase peptide and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the phosphatase peptide.

In some uses, the fusion protein does not affect the activity of the phosphatase peptide *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant phosphatase peptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A phosphatase peptide-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the phosphatase peptide.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the proteins of the present invention, such as naturally occurring mature forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogues of the peptides. Such variants can readily be generated using art-known techniques

in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the phosphatase peptides of the present invention. The degree of homology/identity present will be based primarily on whether the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.

To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of the length of a reference sequence is aligned for comparison purposes. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von

Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated
 5 into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.*
 10 *12(1)*:387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Myers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped
 25 alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

Full-length pre-processed forms, as well as mature processed forms, of proteins that
 30 comprise one of the peptides of the present invention can readily be identified as having complete sequence identity to one of the phosphatase peptides of the present invention as

well as being encoded by the same genetic locus as the phosphatase peptide provided herein. The gene encoding the novel phosphatase protein of the present invention is located on a genome component that has been mapped to human chromosome 2 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data.

5 Allelic variants of a phosphatase peptide can readily be identified as being a human protein having a high degree (significant) of sequence homology/identity to at least a portion of the phosphatase peptide as well as being encoded by the same genetic locus as the phosphatase peptide provided herein. Genetic locus can readily be determined based on the genomic information provided in Figure 3, such as the genomic sequence mapped to the
10 reference human. The gene encoding the novel phosphatase protein of the present invention is located on a genome component that has been mapped to human chromosome 2 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data. As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a phosphatase peptide encoding nucleic acid molecule under stringent conditions as more fully described below.

Figure 3 provides information on SNPs that have been found in the gene encoding the phosphatase protein of the present invention. 21 SNPs were identified. Changes in the amino acid sequence caused by these SNPs can readily be determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference. These SNPs may also affect control/regulatory elements. Positioning of each SNP in an exon, intron, or outside the ORF can readily be determined using the DNA position given for each
20 SNP and the start/stop, exon, and intron genomic coordinates given in Figure 3.

Paralogs of a phosphatase peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the phosphatase peptide, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are
30 typically at least about 60% or greater, and more typically at least about 70% or greater homology through a given region or domain. Such paralogs will be encoded by a nucleic

acid sequence that will hybridize to a phosphatase peptide encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a phosphatase peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the phosphatase peptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a phosphatase peptide encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the phosphatase peptides of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the phosphatase peptide. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those that substitute a given amino acid in a phosphatase peptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant phosphatase peptides can be fully functional or can lack function in one or more activities, e.g. ability to bind substrate, ability to dephosphorylate substrate, ability to mediate signaling, etc. Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Figure 2 provides the result of protein analysis and can be used to identify critical domains/regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 244:1081-1085 (1989)), particularly using the results provided in Figure 2. The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as phosphatase activity or in assays such as an *in vitro* proliferative activity. Sites that are critical for binding partner/substrate binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.* *Science* 255:306-312 (1992)).

The present invention further provides fragments of the phosphatase peptides, in addition to proteins and peptides that comprise and consist of such fragments, particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that may be disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16, or more contiguous amino acid residues from a phosphatase peptide. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the phosphatase peptide or could be chosen for the ability to perform a function, e.g. bind a substrate or act as an immunogen. Particularly important fragments are biologically active fragments, peptides that are, for example, about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the phosphatase peptide, e.g., active site, a transmembrane domain or a substrate-binding domain. Further, possible fragments include, but are not limited to, domain or motif containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well known and readily available to those of skill in the art (e.g., PROSITE analysis). The results of one such analysis are provided in Figure 2.

Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including

the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in phosphatase peptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the phosphatase peptides of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature phosphatase peptide is fused with another compound, such as a compound to increase the half-life of the phosphatase peptide, or in which the additional amino acids are fused to the mature phosphatase peptide, such as a leader or secretory sequence or a sequence for purification of the mature phosphatase peptide or a pro-protein sequence.

Protein/Peptide Uses

The proteins of the present invention can be used in substantial and specific assays related to the functional information provided in the Figures; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its binding partner or ligand) in biological fluids; and as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein or ligand (such as, for example, in a phosphatase-effector protein interaction or phosphatase-ligand interaction), the protein can be used to identify the binding partner/ligand so as to develop a system to identify inhibitors of the binding interaction. Any or all of these uses are capable of being developed into reagent grade or kit format for commercialization as commercial products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

The potential uses of the peptides of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, phosphatases isolated from humans and their human/mammalian orthologs serve as targets for identifying agents for use in mammalian therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the phosphatase. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes. A large percentage of pharmaceutical agents are being developed that modulate the activity of phosphatase proteins, particularly

members of the PPM1G subfamily (see Background of the Invention). The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the expression information provided in Figure 1. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. Such uses can readily be determined using the information provided herein, that which is known in the art, and routine experimentation.

The proteins of the present invention (including variants and fragments that may have been disclosed prior to the present invention) are useful for biological assays related to phosphatases that are related to members of the PPM1G subfamily. Such assays involve any of the known phosphatase functions or activities or properties useful for diagnosis and treatment of phosphatase-related conditions that are specific for the subfamily of phosphatases that the one of the present invention belongs to, particularly in cells and tissues that express the phosphatase. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes.

The proteins of the present invention are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express the phosphatase, as a biopsy or expanded in cell culture. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the phosphatase protein.

The polypeptides can be used to identify compounds that modulate phosphatase activity of the protein in its natural state or an altered form that causes a specific disease or

pathology associated with the phosphatase. Both the phosphatases of the present invention and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to the phosphatase. These compounds can be further screened against a functional phosphatase to determine the effect of the compound on the phosphatase activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the phosphatase to a desired degree.

Further, the proteins of the present invention can be used to screen a compound for the ability to stimulate or inhibit interaction between the phosphatase protein and a molecule that normally interacts with the phosphatase protein, e.g. a substrate or a component of the signal pathway that the phosphatase protein normally interacts (for example, another phosphatase). Such assays typically include the steps of combining the phosphatase protein with a candidate compound under conditions that allow the phosphatase protein, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the phosphatase protein and the target, such as any of the associated effects of signal transduction such as protein phosphorylation, cAMP turnover, and adenylate cyclase activation, etc.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')₂, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries).

One candidate compound is a soluble fragment of the receptor that competes for substrate binding. Other candidate compounds include mutant phosphatases or appropriate fragments containing mutations that affect phosphatase function and thus compete for

substrate. Accordingly, a fragment that competes for substrate, for example with a higher affinity, or a fragment that binds substrate but does not allow release, is encompassed by the invention.

The invention further includes other end point assays to identify compounds that modulate (stimulate or inhibit) phosphatase activity. The assays typically involve an assay of events in the signal transduction pathway that indicate phosphatase activity. Thus, the dephosphorylation of a substrate, activation of a protein, a change in the expression of genes that are up- or down-regulated in response to the phosphatase protein dependent signal cascade can be assayed.

Any of the biological or biochemical functions mediated by the phosphatase can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art or that can be readily identified using the information provided in the Figures, particularly Figure 2. Specifically, a biological function of a cell or tissues that expresses the phosphatase can be assayed. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes.

Binding and/or activating compounds can also be screened by using chimeric phosphatase proteins in which the amino terminal extracellular domain, or parts thereof, the entire transmembrane domain or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by heterologous domains or subregions. For example, a substrate-binding region can be used that interacts with a different substrate than that which is recognized by the native phosphatase. Accordingly, a different set of signal transduction components is available as an end-point assay for activation. This allows for assays to be performed in other than the specific host cell from which the phosphatase is derived.

The proteins of the present invention are also useful in competition binding assays in methods designed to discover compounds that interact with the phosphatase (e.g. binding partners and/or ligands). Thus, a compound is exposed to a phosphatase polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide.

5 Soluble phosphatase polypeptide is also added to the mixture. If the test compound interacts with the soluble phosphatase polypeptide, it decreases the amount of complex formed or activity from the phosphatase target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the phosphatase. Thus, the soluble polypeptide that competes with the target phosphatase region is designed to
10 contain peptide sequences corresponding to the region of interest.

To perform cell free drug screening assays, it is sometimes desirable to immobilize either the phosphatase protein, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

15 Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell
20 lysates (e.g., ³⁵S-labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix,
25 separated by SDS-PAGE, and the level of phosphatase-binding protein found in the bead fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target
30 molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a phosphatase-binding protein and a candidate

compound are incubated in the phosphatase protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the phosphatase protein target molecule, or which are reactive with phosphatase protein and compete with the target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the phosphatases of the present invention can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal or other model system. Such model systems are well known in the art and can readily be employed in this context.

Modulators of phosphatase protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the kinase pathway, by treating cells or tissues that express the phosphatase. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. These methods of treatment include the steps of administering a modulator of phosphatase activity in a pharmaceutical composition to a subject in need of such treatment, the modulator being identified as described herein.

In yet another aspect of the invention, the phosphatase proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with the phosphatase and are involved in phosphatase activity. Such phosphatase-binding proteins are also likely to be involved in the propagation of signals by the phosphatase proteins or phosphatase targets as, for example, downstream elements of a kinase-mediated signaling pathway. Alternatively, such phosphatase-binding proteins are likely to be phosphatase inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a phosphatase protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a phosphatase-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the phosphatase protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a phosphatase-modulating agent, an antisense phosphatase nucleic acid molecule, a phosphatase-specific antibody, or a phosphatase-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

The phosphatase proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanomas, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. The method involves

contacting a biological sample with a compound capable of interacting with the phosphatase protein such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The peptides of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered phosphatase activity in cell-based or cell-free assay, alteration in substrate or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

In vitro techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody or other types of detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-

266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes effects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the phosphatase protein in which one or more of the phosphatase functions in one population is different from those in another population. The peptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other substrate-binding regions that are more or less active in substrate binding, and phosphatase activation. Accordingly, substrate dosage would necessarily be modified to maximize the therapeutic effect within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. Accordingly, methods for treatment include the use of the phosphatase protein or fragments.

Antibodies

The invention also provides antibodies that selectively bind to one of the peptides of the present invention, a protein comprising such a peptide, as well as variants and fragments

thereof. As used herein, an antibody selectively binds a target peptide when it binds the target peptide and does not significantly bind to unrelated proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still selective despite some degree of cross-reactivity.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')₂, and Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, Antibodies, Cold Spring Harbor Press, (1989).

In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the phosphatase proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or phosphatase/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence alignment can be used to identify conserved and unique sequence fragments.

An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such fragments can be selected on a physical property, such as

fragments correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2).

Detection on an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

Antibody Uses

The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues in an organism and over the course of normal development. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Also, such antibodies can be used to assess abnormal tissue distribution or abnormal expression during development or progression of a biological condition.

Antibody detection of circulating fragments of the full length protein can be used to identify turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function. When a disorder is caused by an inappropriate tissue distribution, developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. If a disorder is characterized by a specific mutation in the protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the protein or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. Thus, where a specific

protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the phosphatase peptide to a binding partner such as a substrate. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the peptides activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount of protein in the sample; means for comparing the amount of protein in the sample with a standard; and instructions for use. Such a kit can be supplied to detect a single protein or epitope or can be configured to detect one of a multitude of epitopes, such as in an antibody detection array. Arrays are described in detail below for nucleic acid arrays and similar methods have been developed for antibody arrays.

Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a phosphatase peptide or protein of the present invention (cDNA, transcript and genomic sequence). Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide sequence that encodes one of the phosphatase peptides of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.

As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide

sequences, for example up to about 5KB, 4KB, 3KB, 2KB, or 1KB or less, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations described herein such as recombinant expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a transcript/cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide sequence is the complete nucleotide sequence of the nucleic acid molecule.

The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists essentially of a nucleotide sequence when such a nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ

ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional nucleotides or can comprises several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, humans genomic sequence (Figure 3) and cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when the mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the phosphatase peptide alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and

non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

5 Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

10 The invention further provides nucleic acid molecules that encode fragments of the peptides of the present invention as well as nucleic acid molecules that encode obvious variants of the phosphatase proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

15 The present invention further provides non-coding fragments of the nucleic acid molecules provided in Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences and gene termination sequences. Such fragments are useful in controlling heterologous gene expression and in developing screens to identify gene-modulating agents. A promoter can readily be identified as being 5' to the ATG start site in the genomic sequence provided in Figure 3.

25 A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could at least 30, 40, 50, 100, 250 or 500 nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope bearing regions of the peptide, or can be useful as DNA probes

and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further, primers can be used in PCR reactions to clone specific regions of gene.

5 A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 20, 25, 40, 50 or more consecutive nucleotides.

10 Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. Allelic variants can readily be determined by genetic locus of the encoding gene. The gene encoding the novel phosphatase protein of the present invention is located on a genome component that has been mapped to human chromosome 2 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data.

20 Figure 3 provides information on SNPs that have been found in the gene encoding the phosphatase protein of the present invention. 21 SNPs were identified. Changes in the amino acid sequence caused by these SNPs can readily be determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference. These SNPs may also affect control/regulatory elements. Positioning of each SNP in an exon, intron, or outside the ORF can readily be determined using the DNA position given for each SNP and the start/stop, exon, and intron genomic coordinates given in Figure 3.

25 As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The conditions can be such that sequences at least about 60%, at least about 30 70%, or at least about 80% or more homologous to each other typically remain hybridized to

each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6.

One example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65C. Examples of moderate to low stringency hybridization conditions are well known in the art.

Nucleic Acid Molecule Uses

The nucleic acid molecules of the present invention are useful for probes, primers, chemical intermediates, and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the same or related peptides shown in Figure 2. As illustrated in Figure 3, 21 SNPs were identified.

The probe can correspond to any sequence along the entire length of the nucleic acid molecules provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be construed as encompassing fragments disclosed prior to the present invention.

The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.

The nucleic acid molecules are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

The nucleic acid molecules are also useful as probes for determining the chromosomal positions of the nucleic acid molecules by means of *in situ* hybridization methods. The gene encoding the novel phosphatase protein of the present invention is located on a genome component that has been mapped to human chromosome 2 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data.

The nucleic acid molecules are also useful in making vectors containing the gene regulatory regions of the nucleic acid molecules of the present invention.

The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful as hybridization probes for determining the presence, level, form and distribution of nucleic acid expression. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes. Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic acid whose level is determined can be DNA or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or decrease in phosphatase protein expression relative to normal results.

In vitro techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA includes Southern hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a phosphatase protein, such as by measuring a level of a phosphatase-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if a phosphatase gene has been mutated. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes.

Nucleic acid expression assays are useful for drug screening to identify compounds that modulate phosphatase nucleic acid expression.

The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the phosphatase gene, particularly biological and pathological processes that are mediated by the phosphatase in cells and tissues that express it. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes. The method typically includes assaying the ability of the compound to modulate the expression of the phosphatase nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired phosphatase nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the phosphatase nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

The assay for phosphatase nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds involved in the signal pathway. Further, the expression of genes that are up- or down-regulated in response to the phosphatase protein signal pathway can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of phosphatase gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of phosphatase mRNA in the presence of the candidate compound is compared to the level of expression of phosphatase mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate phosphatase nucleic acid expression in cells and tissues that express the phosphatase. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or nucleic acid expression.

Alternatively, a modulator for phosphatase nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits the phosphatase nucleic acid expression in the cells and tissues that express the protein. Experimental data as provided in Figure 1 indicates expression in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, lung small cell carcinomas, and leukocytes.

The nucleic acid molecules are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of the phosphatase gene in clinical

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in phosphatase nucleic acid expression, and particularly in qualitative changes that lead to pathology. The nucleic acid molecules can be used to detect mutations in phosphatase genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally occurring genetic mutations in the phosphatase gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of the phosphatase gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a phosphatase protein.

38

indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data. Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364 (1994)), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.*, *Nucleic Acids Res.* 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

Alternatively, mutations in a phosphatase gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant phosphatase gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve, C.W., (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International

Publication No. WO 94/16101; Cohen *et al.*, *Adv. Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144 (1993); and Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include selective oligonucleotide hybridization, selective amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship). Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the phosphatase gene in an individual in order to select an appropriate compound or dosage regimen for treatment. Figure 3 provides information on SNPs that have been found in the gene encoding the phosphatase protein of the present invention. 21 SNPs were identified. Changes in the amino acid sequence caused by these SNPs can readily be determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference. These SNPs may also affect control/regulatory elements. Positioning of each SNP in an exon, intron, or outside the ORF can readily be determined using the DNA position given for each SNP and the start/stop, exon, and intron genomic coordinates given in Figure 3.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual.

Accordingly, the production of recombinant cells and animals containing these

polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control phosphatase gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of phosphatase protein. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into phosphatase protein.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of phosphatase nucleic acid. Accordingly, these molecules can treat a disorder characterized by abnormal or undesired phosphatase nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the phosphatase protein, such as substrate binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in phosphatase gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired phosphatase protein to treat the individual.

The invention also encompasses kits for detecting the presence of a phosphatase nucleic acid in a biological sample. Experimental data as provided in Figure 1 indicates that phosphatase proteins of the present invention are expressed in humans in the brain, breast, colon adenocarcinomas, prostate, skin melanotic melanoms, eye retinoblastomas, placenta choriocarcinomas, uterus leiomyosarcomas, and lung small cell carcinomas, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in human leukocytes. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting phosphatase nucleic acid in a biological sample; means for determining the amount of phosphatase nucleic acid in the sample; and means for comparing the amount of phosphatase nucleic acid in the sample

with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect phosphatase protein mRNA or DNA.

5 Nucleic Acid Arrays

The present invention further provides nucleic acid detection kits, such as arrays or microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1 and 3).

As used herein "Arrays" or "Microarrays" refers to an array of distinct
 10 polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared and used according to the methods described in US Patent 5,837,832, Chee *et al.*, PCT application W095/11995 (Chee *et al.*), Lockhart, D. J. *et al.* (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. *et al.* (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their
 15 entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown *et al.*, US Patent No. 5,807,522.

The microarray or detection kit is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense
 20 oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray or detection kit, it may be preferable to use oligonucleotides that are only 7-20 nucleotides in length. The microarray or detection kit may contain oligonucleotides
 25 that cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray or detection kit may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray or
 30 detection kit, the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm which starts at the 5' or at

the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray or detection kit. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler *et al.*) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray or detection kit, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray or detection kit so that the probe sequences hybridize to complementary oligonucleotides of the microarray or detection kit. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each

oligonucleotide sequence on the microarray or detection kit. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large-scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression of the phosphatase proteins/peptides of the present invention. In detail, such methods comprise incubating a test sample with one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention and or alleles of the phosphatase gene of the present invention. Figure 3 provides information on SNPs that have been found in the gene encoding the phosphatase protein of the present invention. 21 SNPs were identified. Changes in the amino acid sequence caused by these SNPs can readily be determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference. These SNPs may also affect control/regulatory elements. Positioning of each SNP in an exon, intron, or outside the ORF can readily be determined using the DNA position given for each SNP and the start/stop, exon, and intron genomic coordinates given in Figure 3.

Conditions for incubating a nucleic acid molecule with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the Human genome disclosed herein. Examples of such assays can be found in Chard, T, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of Enzyme Immunoassays*:

Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the Human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid.

In detail, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified phosphatase gene of the present invention can be routinely identified using the sequence information disclosed herein can be readily incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in prokaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the nucleic acid molecules from the vector.

Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage λ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

5 In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences
10 that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example
15 vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and
20 eukaryotic hosts are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types
25 such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is
30 joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures

for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques.

5 Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the
10 peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa,
15 thrombin, and enterophosphatase. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, *Gene* 69:301-
20 315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)).

Recombinant protein expression can be maximized in host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in*
25 *Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are
30 operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-

943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the nucleic acid molecule sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill

in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989*).

Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

Where secretion of the peptide is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as phosphatases, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

Where the peptide is not secreted into the medium, which is typically the case with phosphatases, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the peptides described herein, the peptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may include an initial modified methionine in some cases as a result of a host-mediated process.

Uses of vectors and host cells

The recombinant host cells expressing the peptides described herein have a variety of uses. First, the cells are useful for producing a phosphatase protein or peptide that can be further purified to produce desired amounts of phosphatase protein or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the phosphatase protein or phosphatase protein fragments, such as those described above as well as other formats known in the art. Thus, a recombinant host cell expressing a native phosphatase protein is useful for assaying compounds that stimulate or inhibit phosphatase protein function.

Host cells are also useful for identifying phosphatase protein mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant phosphatase protein (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native phosphatase protein.

Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such

as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a phosphatase protein and identifying and evaluating modulators of phosphatase protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the phosphatase protein nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the phosphatase protein to particular cells.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example

of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al. PNAS* 89:6232-6236 (1992).

Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al. Science* 251:1351-1355 (1991). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein are useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect substrate binding, kinase protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* phosphatase protein function, including substrate interaction, the effect of specific mutant phosphatase proteins on phosphatase protein function and substrate interaction, and the effect of chimeric phosphatase proteins. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more phosphatase protein functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method

and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

SEQUENCE LISTING

<110> WEI, Ming-Hui et al

<120> ISOLATED HUMAN PHOSPHATASE PROTEINS,
 NUCLEIC ACID MOLECULES ENCODING HUMAN PHOSPHATASE PROTEINS,
 AND USES THEREOF

<130> CL001066-CIP

<150> 09/752,820

<151> 2001-01-03

<160> 7

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1435

<212> DNA

<213> Human

<400> 1

```

tcgcgccttt caccggcacc ttgcgtcggt cgcgccgcgg ggcttgcctc tgccgcgcgc 60
acccccgggg ctteggctcc ggcacgggtc ggcgccagct ttctgcacc tgaggccgcc 120
ggccagccgc cgccatgggt gcctacctct cccagcccaa cacggtgaag tgctccgggg 180
acggggtcgg cgcgccgcgc ctgccgtgc cctacggctt ctccgccatg caaggctggc 240
gcgtctccat ggaggatgct cacaactgta ttctgagct ggacagtga acagccatgt 300
tttctgtcta cgatggacat ggaggggagg aagttgcctt gtactgtgcc aaatatcttc 360
ctgatatcat caaagatcag aaggcctaca aggaaggcaa gctacagaag gctttagaag 420
atgccttctt ggctattgac gccaaattga ccactgaaga agtcattaaa gagctggcac 480
agattgcagg gcgacccact gaggatgaag atgaaaaaga aaaagtagct gatgaagatg 540
atggggacca cttctataag agaaacaaga acctgccacc tgaggaaacag atgatttcag 600
cccttcctga catcaagggt ctgactctca ctgacgacca tgaattcatg gtcattgcct 660
gtgatggcat ctggaatgtg atgagcagcc aggaagttgt agatttcatt caatcaaaga 720
tcagccagcg tgatgaaaat ggggagcttc ggttattgtc atccattgtg gaagagctgc 780
tgatcagtg cctggcacca gacacttctg gggatggtag aggggtgtgac aacatgacct 840
gcatcatcat ttgcttcaag ccccgaaaca cagcagagct ccagccagag agtggcaagc 900
gaaaactaga ggaggtgctc tctactgagg gggctgaaga aaatggcaac agcgacaaga 960
agaagaaggc caagcgagac tagcagtcac ccagaccctt gccacctag actgttttct 1020
gagccctccg gacctgagac tgagttttgt ctttttcctt tagccttagc agtgggtatg 1080
agggtgtgcag ggggagctgg gtggcttcac tccgcccatt ccaaagaggg ctctccctcc 1140
acactgcagc cgggagcctc tgctgtcctt cccagccgcc tctgctcctc gggctcatca 1200
ccggttctgt gcctgtgctc tgttgtgttg gagggaagga ctggcgggtc tgggtttttac 1260
tctgtgaact ttatttaagg acattctttt ttattggcgg ctccatggcc ctggccgct 1320
tgcacccgct ctctgttgta cactttcaat caacactttt tcagactaaa ggccaaaacc 1380
taatcgttaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 1435

```

<210> 2

<211> 282

<212> PRT

<213> Human

<400> 2

Met Gly Ala Tyr Leu Ser Gln Pro Asn Thr Val Lys Cys Ser Gly Asp

1	5	10	15
Gly Val Gly Ala Pro Arg Leu Pro Leu Pro Tyr Gly Phe Ser Ala Met			
	20	25	30
Gln Gly Trp Arg Val Ser Met Glu Asp Ala His Asn Cys Ile Pro Glu			
	35	40	45
Leu Asp Ser Glu Thr Ala Met Phe Ser Val Tyr Asp Gly His Gly Gly			
	50	55	60
Glu Glu Val Ala Leu Tyr Cys Ala Lys Tyr Leu Pro Asp Ile Ile Lys			
	65	70	75
Asp Gln Lys Ala Tyr Lys Glu Gly Lys Leu Gln Lys Ala Leu Glu Asp			
	85	90	95
Ala Phe Leu Ala Ile Asp Ala Lys Leu Thr Thr Glu Glu Val Ile Lys			
	100	105	110
Glu Leu Ala Gln Ile Ala Gly Arg Pro Thr Glu Asp Glu Asp Glu Lys			
	115	120	125
Glu Lys Val Ala Asp Glu Asp Asp Gly Asp His Phe Tyr Lys Arg Asn			
	130	135	140
Lys Asn Leu Pro Pro Glu Glu Gln Met Ile Ser Ala Leu Pro Asp Ile			
	145	150	155
Lys Val Leu Thr Leu Thr Asp Asp His Glu Phe Met Val Ile Ala Cys			
	165	170	175
Asp Gly Ile Trp Asn Val Met Ser Ser Gln Glu Val Val Asp Phe Ile			
	180	185	190
Gln Ser Lys Ile Ser Gln Arg Asp Glu Asn Gly Glu Leu Arg Leu Leu			
	195	200	205
Ser Ser Ile Val Glu Glu Leu Leu Asp Gln Cys Leu Ala Pro Asp Thr			
	210	215	220
Ser Gly Asp Gly Thr Gly Cys Asp Asn Met Thr Cys Ile Ile Ile Cys			
	225	230	235
Phe Lys Pro Arg Asn Thr Ala Glu Leu Gln Pro Glu Ser Gly Lys Arg			
	245	250	255
Lys Leu Glu Glu Val Leu Ser Thr Glu Gly Ala Glu Glu Asn Gly Asn			
	260	265	270
Ser Asp Lys Lys Lys Lys Ala Lys Arg Asp			
	275	280	

<210> 3
 <211> 29695
 <212> DNA
 <213> Human

<220>
 <221> misc_feature
 <222> (1)...(29695)
 <223> n = A,T,C or G

<400> 3
 aaagaatcctt tttttttttt ttgagacgga gttgctctgt caccaggggt ggagtgcagt 60
 ggcgccatct tggttcactg caacctccgc ctctgggggt caagctattc gcctgcctta 120
 gcctcccaag tagctgggat tacaggagcg caccactacg cctggctaata ttttgtattt 180
 ttagtagaga cgggtttcac atgttgcca ggctgggtct gaacttctgg cctcaagtga 240
 tccaccaccc ccccttgcc tcccaaagt ctgggattac aagtgtgagc cactgtgccc 300
 ggctgaaaag aatcaatttt gtcatagttt ggagaatttc tccttttctc tccatccctt 360
 gaatgcaatt tattaccaaa tctgtcttat ttgttattgt ctaatttgct ctttcatctg 420
 gattcccatt gccaccctgc gtggtaccac cttactccca gctcttctca tctcctgctt 480
 agagtaagag ctctctaact agtagcagtg cccagggcca ggcgcggtgg ctcacgcctg 540

taatcccagc	acttttgggag	gctgagggcgg	gtagatcacg	aggtcaggag	ttcgagacca	600
gcctggccaa	catggtgaaa	ccccgtctct	actaaaaata	caaaaattag	ccaggcgtgg	660
aatcctagct	actcgggagg	ctgagggcagg	agaatcgctt	gaacctggga	ggcggagggt	720
gcagtgagcc	aagatagcgc	cactgcactg	cagcctgggc	aacaagagcg	aaactctgtc	780
ttaaaaaaaaa	aacaatagta	ggcgggggtgc	ggtgggtcac	gcctgtaatc	ccaacacttt	840
gggaggccga	ggcggggcgga	tcacgaggtc	aggagatgga	gaccatcctg	gctaacacgg	900
tgaaacctcg	tctctactaa	aaatacaaaa	aattagccag	gcgtgggtggc	gggcgcctgt	960
agtcccagct	actcgggagg	ctgagggcagg	agaatggcgt	gaacccggga	ggcggagcct	1020
gtagcctggg	cgacagagcg	agactctgtc	tcaaaaacaa	acaaaaaaca	acaacacaac	1080
agtgcctcag	actctctccc	tccaatgtac	actgcataca	aagactagac	aaacgatgcc	1140
aaagggttaca	ccaggacgac	aatgaagtcc	aagtcactca	tcttggcgct	tctttttatc	1200
aagctagcta	gttaatatata	ccacttacaa	tcattctcctg	agtccagcca	gcttcctcag	1260
aaatcctccc	caaatgcagt	tcacattctt	ccctcactcc	tccaaacctt	ataacattat	1320
tccctttttcc	tgtggcataa	tgcaatccag	aggcatcctc	tctgagagcc	tacatccttt	1380
caatcctcca	agaaacagct	cctcctccca	tccttgagct	tttcccaccc	agaataggct	1440
gtaccaaaca	tttctacatt	gtattaactc	gaattatata	acagatcgag	gtcttcgctt	1500
tcagaaaaga	ctcacattct	tctcatatag	gcctcgcctg	gcctggccaa	gtacttccta	1560
agctcggaac	aaatactggt	caactttaat	tgaaccaa	cgggcggggt	ttgcggagtc	1620
tgagagtgc	acgttgggga	gagggggatg	aaaacactgg	aggacgggct	gaaagcgctc	1680
agtcgcgac	aaaagaggcg	tcagacaaaa	cgccaagagg	ctggggactg	ggaacgaagg	1740
aagaggttct	gccagaggcg	acctgccacc	tgcgcgagga	agcggagtag	gacggcgggc	1800
gttgggtggc	gtggtcgcgc	tagtctcgcg	ggagcggccg	ttgggcgggc	cgttgtcccc	1860
tgcgggcggg	gcgagttgct	aaggaaatga	ctgcccgcag	cgcctggccc	cgccgcgcag	1920
gccgggcggg	gtctggagcg	gcgccttttc	cgcctccgct	ccctcacagc	tcccgctccc	1980
ttaccgcctc	ctggccggcc	tcgcgccttt	caccggcacc	ttgcgtcggg	cgcgcgcgcg	2040
ggcctgctcc	tgccgcgcgc	acccccgggg	cttcggctcc	ggcacgggtc	gcgcccagct	2100
ttcctgcacc	tgagccgcgc	gccagccgcc	gccatgggtg	cctacctctc	ccagcccaac	2160
acggtgaagt	gctccgggga	cggggctcgc	gccccgcgcc	tgccgctgcc	ctacggcttc	2220
tccgccatgc	aaggctggcg	cgtctccatg	gaggtgagga	ggcaggggcc	cataggctgg	2280
ccgctgcggg	gcgggaatct	gacggagaaa	gagagcgggg	gatgggggtc	tcccctggga	2340
aggggtccaa	ttgggagcct	gcggccgcag	cggccgtttg	cggggcgaca	gagaccgcgg	2400
ggtcagggcc	gagagggagc	tcccgatgct	tggggaccgt	gtgccgggtg	ccagtggcgg	2460
cgagggctcg	gccatgtggg	aagaggcacc	ttccgcccac	tgaccgccct	ctccccgagc	2520
tttggcgcca	tcctctcgtg	ccaacctagc	cctccaggct	catcaacggt	gtgagggtta	2580
gtgtgggagt	aaagacgcaa	ataggggcta	tttattcatt	tgtttttcaa	gggagagggt	2640
aatagatgtg	aataactttt	taaattttta	tatttaaaat	atctgatgtg	ggaagcctct	2700
tttggctagg	agtttgacag	tgaaggaac	cccgggcaga	gtctgtttca	catttttggt	2760
gcctggcctt	gggctcttgc	atgttaattt	cagaggctgg	acccgacctc	caggagtgtg	2820
cactcatttg	cactcttttc	agggcctttt	actaacttcg	gaaaacttga	attatgtcag	2880
tccttaggtt	ttccttttta	ttatgcttcg	ttttctttct	ctcttaggat	ttctctaaaa	2940
cttaatcagt	aattctcatt	tgctgtaat	tgtagtattat	agtttctttg	attgtaagtc	3000
atttcatcaa	tttttttcat	cacaacctac	ctacaaaggg	cttttctaga	aaattttact	3060
ctggacaaaa	ggggaaaaga	aaaatattgg	ggggaaagta	gtagtattag	gtaaaacttg	3120
atgtgaaact	acaaaagaga	agagggaaaa	ctgcggtagg	gaggaaaggg	aggaagacgg	3180
gttaaccgtg	gctttgtgaa	gagcattctg	aagtctaggc	aaaagggcc	gggaaatact	3240
ctgtctggta	ttgagggttt	ctccacctac	cgggtgggct	tcaggtaaca	gcgaaatact	3300
gtctcccttg	ggaattgttt	cagatccctc	gctcctcctg	tggttagctc	tggaatgcca	3360
gtaggaactt	caatgttttg	ttttccgatt	caaattttat	attcataact	gaccttaata	3420
acaattttac	aattaggtat	aaaattttcag	gatactagtg	tatcctatag	ttcatctcat	3480
ctgctttggc	tccctttttt	tttttttgag	acagaatttc	gctcttggtg	cccaggctgg	3540
aatgcaatgg	cgcgatctcg	gctcaccgca	acctccgcct	cccagggttg	agcaattctc	3600
ctgcctcagc	ctccatagta	gttgggatca	caggcatgtg	ccaccacgcg	tggttatttt	3660
tgtattttta	gtagagacag	ggtttcttca	tgtgggtcag	gctgggtgtg	aactcctgac	3720
ttaggtgatc	tgctgcctc	ggtctcccaa	agtgtacca	ccacnnnnnn	nnnnnnnnnn	3780
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	3840
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	3900
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	3960

nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4020
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4080
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4140
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4200
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4260
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4320
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4380
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4440
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4500
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4560
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4620
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4680
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4740
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4800
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4860
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4920
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4980
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5040
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5100
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5160
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5220
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5280
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5340
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5400
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5460
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5520
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5580
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5640
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5700
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5760
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5820
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5880
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	5940
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	6000
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	6060
nnnnnnnnnn	nnnnntggga	ttacagacat	gagccactgc	gccagcctt	atttagaaat	6120
ttcttttagtg	aaagatgata	aattttcagt	ttttcattat	ctgaacatgt	ttttatctag	6180
cctttgttct	gaaaagatgc	ttggactcag	taccagttc	tagattgaca	gttaattttt	6240
cttaatttgt	aatgttgtt	tcattgattg	acttccattg	ttgttcggaa	aaatttatca	6300
tcagccattt	ctgacttttg	atctgtgttt	tctcttttgt	ttctcttttt	ttttttcttt	6360
tttttttttt	gagacggagt	gtcgtctgt	tgccaggtt	ggagtgcagt	ggcatgatct	6420
tggtcactg	caacctctgc	ctcctgagtt	caagcgattc	tcctgcctca	gcttcccag	6480
tagctgggat	tacagggccc	tgccaccatg	cccggctaata	tttttgtatt	ttcaatagag	6540
acagggtttc	actatgttgg	ccgggttggt	ctccaactcc	tgacctctta	atccgcccgc	6600
ctcggcctcc	caaagtgtcg	gcattacagg	cgtgagccac	catgcctggc	ccatcagttg	6660
atgtagtctt	aaggggacaa	gagtacattt	aatatattgt	tgtgaggttc	tctggaagtg	6720
acaaaactgc	tttctatgga	gagttaggaa	tttttttttt	ttttgaaacg	gagtctcgca	6780
ttgtcaccgc	ggctggagtg	tagtggcttg	atctcggctc	actgaaactt	ccgectcttg	6840
ggttcaagtg	attctcctgc	ctcagccttc	caaatagtcg	ggattacagg	agtctgccac	6900
caggccagct	aatttttttt	tgtattttta	gtggagacag	gatttcacta	tgttggccag	6960
gctggtctca	agactcctga	cgttgtgatc	cacctgcctc	ggcctcccaa	agtgtctgga	7020
ttacagatgt	gagctaccgt	gcccggccag	gaattttttg	tgctataaat	catattttcc	7080
ttttattaaa	ggcagtgatc	atatctatag	tataattttg	aggaggctgg	ctattttattg	7140
ctgtgtagaa	gctggcttat	tagtgggtcaa	ggggtcatct	agaattgact	ataaagatag	7200
tattgagcag	aaaattctta	aaatacctgc	atattagttt	cagtcattaa	attaatggaa	7260
aaaatataaa	aagaaatatc	acaagtatgc	tatgggggtc	taccttaggg	cattgaaggt	7320
tgaaaaacat	ttattttctt	atcttcagat	tagcatctca	tatcaatagc	caatagctta	7380

aagcgcctttt	tacttactaa	accaggtcag	aatttctctc	tctctctctt	tttttttttt	7440
agacagagtc	tctgtcacc	agtctagagt	gcagtggtgt	gatctcggct	cactggcctt	7500
tctgtttaag	tgattctccg	gcctcagcct	cctgagtagc	tgggattaca	ggcatgtgcc	7560
acgacacttg	gcttattttt	tgtattttta	gtagagatgg	ggtttctctg	tgttggtcag	7620
gcaggtcttg	aactcctgac	ctcaggtgat	ctgcccgcct	ccgcctccca	aagtgcctgg	7680
attacagacg	tgagctactg	cgcctggcca	gaatttcttt	gtctagaatg	tggttagcaa	7740
cttttataaa	aacgcattat	ttgcatttga	ttagcatgca	gtaccatttc	acagttcaaa	7800
gctagtatat	aattatatca	catgtatgcc	catgagcatg	gagaaactat	tttcttttta	7860
tttttttaag	ttggagtttt	gctcttggtg	cccaggctgg	agtgcattgg	tgccatctcg	7920
gctcactgca	gcttctgcct	cctgggttct	agcaatttgc	ctaccccagc	ctcccaagta	7980
gctgggatta	caggcactcg	ccaccatgcc	cagctaactt	ttttgtattt	ttagtagaga	8040
aggggttttc	ccatgctggc	gagtctgggt	ttgaacttct	ggcctcaagt	gatctgcccg	8100
cctcagcctc	ccaaagtgtc	ggaattacag	gcatgagcca	ctgtgcccgg	cctttttatt	8160
ttttaaaatta	tttatgtatt	tatttttgaga	caggatctca	ctcttgccca	tgcttgagtg	8220
gtaaggagta	tgggatttct	tgtgccgggt	ccttcacatc	ctcactacac	ttatctgcct	8280
ttacagtggc	tcgatcatag	ttcactgcat	agccttctgg	gctcaagggg	tcttccagcc	8340
tcagcctaata	ataggcacat	gccaccatga	ctggctaatt	tttttttttt	aagttttttt	8400
ttgtagagat	agggccttgc	agtgttgccc	aggctgagga	atttttattt	tgtttatttt	8460
atttatttat	ttattttatt	atttatttat	ttattttatt	attgagatgg	agtcttacac	8520
tgtcaccag	gctggagtgc	agttgcgcgg	tctcagctac	tgcaagctcc	gcctcccggg	8580
ttcatgccat	tctcctgcct	cagcctcccg	cgtagctgga	ctacaggcgc	ccgccaccac	8640
gcctggctaa	tttttttgtgt	gtgttttttag	tagaggcggg	gtttcaccat	gttagccagg	8700
atgggtctga	ttcctgacct	cgtgatccac	ccacatcggc	ctcccaaagt	gctgggatta	8760
caggtgtgag	ccaccatgcc	tgggtctagaa	attattttat	atttttatac	attgccttat	8820
aagttctcaa	gcaactggaa	aatacaatca	gaacgtattc	ctcaagattt	caaggatatt	8880
ttacacaaaag	ttctattgtc	tgattcctta	gcagttgtta	ctactgtttc	cctaacctct	8940
aatcttctat	tgggttatta	gtcttagaat	tgaattttga	gaggtaaggg	cttgaatttg	9000
aacatagaaa	tttatacagg	tctgatcagt	agttcttgac	attgtattat	ctggaaacaa	9060
atcttttagaa	ctgagcttaa	gatgtttaat	gacattttgt	agacagagta	tgatttcagt	9120
gtagttgttt	ttgtttcttt	ctagatctag	ttcagagatg	aagtatatca	actttttttt	9180
cctttttgac	ccaatgctag	cagaaaaaca	acacctttta	atcataattt	gtatttgaaa	9240
atgtgtatac	aggttccttt	ttattttatt	tatttcttac	aggttccttt	ttaatcagct	9300
ttattgagat	agagttcata	tactgtatgg	ttcataccac	atatgggttc	tataccatac	9360
agtatatgaa	ctcactttaa	gagtataatt	cagtgggttt	aagggtataa	ttcattcatt	9420
ttaagggtat	aattcagtgg	cttttagtat	attttctttt	ttttcttctt	tttttctttt	9480
ttcgagacag	ggtcttgctc	tgttacccag	gctggagtgc	agtgcgcag	cctcagctca	9540
ctgcaacctc	cacctccag	gttcaagcga	attctcttgc	ctcagcatcc	tgagttagctg	9600
gaactacagg	ctcacgccac	cacaccagc	taatttttat	attttcagta	gaaacagggt	9660
ttcaccatgt	tgcccagggt	ggtctcgaac	tcctggcctc	aagtgatctg	cctgcctcag	9720
cctcccaaag	tgtctgggatt	acaggcatga	gccaccgtgc	ctgcctgttt	tgtagtgtat	9780
tcaaacagag	ttgtacaact	gtcaccacaa	tcagtttttag	aaccccaaaa	agaaacctg	9840
tactctttac	cagtcactcc	ctatcttccg	tcactaacc	cctggcatcc	actaatttac	9900
atgacctcta	tgaatttgcc	tattctgaac	attttataaa	tggatttcta	aatacactac	9960
cttttatatac	tggatgcttt	tactaagcat	gtgtattttt	gaaattgact	ttaaagcttg	10020
ttggcccttg	gaagagtaaa	ttactctcca	ccccaagtat	tcctctacc	cctcagcttt	10080
gcctgtaagt	ttctttttta	aaaaatcaca	catacattgt	tgtagtagat	ttagaataa	10140
gtattttgct	gacccaaggt	tcttttgctt	ctttctagat	cagtgccttg	caggttttat	10200
tttacagagc	ttaatagaat	cagaaatctc	tttaaaaactc	cagtctcata	tccagttatc	10260
actcaccatc	tctgtgtttg	cagcaatagc	caggcctggc	ccagagggac	ttgatctcca	10320
cttttggttt	ttagactttt	ctgtggcttt	taccacctgc	tgtgtatcct	tgacctcata	10380
ctgctggact	cctttggatg	gataccagca	ggatgggttc	ggctccagtg	ggcacttttt	10440
aaaattctct	ccttctgttc	agatagacag	agctcaggca	gatcaccaag	tctgttgccct	10500
gtgtaaccag	gaaagagatg	ctagttttct	tttaggcact	cccatttggt	tctgttgcaa	10560
ccttcctcac	ttagttgatg	gaagggaagc	aaaagaccca	gaactccatc	tcaaattaat	10620
gacttaacaa	ttcttgaatt	ttctcttata	tcctagttaa	ctctttttct	tatctccagg	10680
aggtcagttt	taattattgt	tgtttattca	tttattttca	tggagacagg	gtcttactgt	10740
gttgtccagg	ctggccttga	accctgggcc	tcaaggaatc	ttttcacctc	agcctcccaa	10800


```

agtgctggaa ttacagggcat gagtcaccac acccagcctg atatttttca gttgatgtat 10860
catagtgtgtg cctaagcata atttttttaat tttaattttt tatttttggg gacaggggtct 10920
ccctctgtcg cccagggctgg agtgcagtga tgcgatctca gctcactgca acctccacct 10980
tctgggttaa agcgattctc ctgcctcagc ctcccagagta gctgggacta caggtaccca 11040
ccatcacacc cggctaaatt ttttgtgtgt atttttagtag agacgagatt tcgctgtgtt 11100
gcccagggcg gtctcgaact cctgagctca ggcaatccgc ctgcctcggc ctcccaaagt 11160
gctgggatta caggcatgaa ccaccacgcc cggccaagcg taatatTTTT aagggtcac 11220
aatgttgtgt catgaatcaa tcagtgtttc gttctttttt atgggtgaat aatattccat 11280
ggatatggatt tgtcacattt tgtttatcca ttcatagatt gatagacatt ttggatttcc 11340
actttttttt tttttttttt gctattataa atagtgtac tatgtacaaa ttttttgtgtg 11400
gaaatatgtc ctcatatctc ttgggtatat accaaagagt ggaagtgtg ggtcatatgg 11460
taactacgtg tttaacattt tgagaaactg ctaaactgtt ttccaaagt gctgtaccgt 11520
cgtacattcc tgccagcaat atatgaggat gccagttcct tcacatgttc actacactta 11580
tccacctttt ttataataac taatgggtggg tgtgagatgg tatctcattg tagttttgat 11640
ttgtatttct ctgatggcta aatgggcta atgtgttgaa ctttttgttt gagacagaat 11700
ctcactctgt ccagattcaa gcgattctcc tgcctcagcc tccctagcag ctgggattac 11760
aggcacatgc caccacaccc agctaatttt ttgtatttct agtagagaca gggcattacc 11820
atgttgttca ggctgggtgc gaactcctga cctcaaagga tccgcctccc tgggcctccc 11880
aaagtgtctg attacaggct agagccacca tgccaggcct tatgtttgaa catcttttat 11940
gtgcttattg gacatttgtg tatcttcttt ggagaaatgt ctgttcaaag tctttgtcca 12000
tttttaattg gattgtcttt ttgtcttttg atgtgtaaga gttctttatg tgttttgat 12060
acaagtttgt tagatatatg atttgcaaat cttttctcca attttttgtg acttttgctt 12120
tctttttttg ttttgttttt gttgttgttg ttgttgttgt tgttttggtc gggggacagt 12180
cttgcctctga ccaccagggc tggaatggag tggcgcgatc ttgggttcaact gcaacctctg 12240
cctcctgagt tcaagctatc ctgcttcagc ctcccagagta gctgggaccc aggtgtgtgc 12300
caccactccc agctaatttt ttatttttag tagagaccgg gtttcacat gttggccagg 12360
ctgggtcttg actactgacc tcaggtgatc tgcctgcctc agcctcccaa agtgctggga 12420
ttacagtcac gagccactac acctgatcc tttttgctgg ctttctttct tttttttct 12480
tttttttttt gagacggagt ctgcctctgt tgccaggctg gagtgcagtg gcatgatctc 12540
ggctcactgc aacctctgcc tcccggttc aagccattct cctgcctcag cctcccgagt 12600
agctaggact ataggcacat gccaccatgc gcagctaatt ttgtatttt tagtagagac 12660
ggggtttcac catgttgccc aggatagtct cgatctcttg acctcgtgat ccgtccgcct 12720
gggcctccca aagtgttagg attacagacg tgagccacca cactcagcct ctttttgctt 12780
tcttgatggt gtcttttgaa acaaaagtgt ttacttttga taaagtccaa tttgtctatt 12840
ttgtttgttt gtttttgtta agaagctttg cctaacccaa agtcacgaga attttctctt 12900
aggttttctt ctaagagttt tatagtttta gctgtttctg tgatccattt tgagtgaatt 12960
tttgtgaatg gtatgaggga gtgatccaac ttcatctctt tgtgtgtgga tatcaagttg 13020
tcccagcact atttgtttaa accactgttc ttttcccca ttgaattatc ttggcatcat 13080
tgtcagagat aaattgaccg taaatgtgag ggttttatct ctgaactctc aagtccattt 13140
cattgggtcta catgtcccta tgccagtaat acactatctt ggttactgta gcttttttagt 13200
acgttttgaa atgtttttta aatttgtttt tcatctaaat tttaggatta atttgtcaat 13260
ttctgcacaa aaggcacctg ggtttctata ggggttatgc agaactctga gatcaactgg 13320
gggagtatta caggcatgag ccaccgtgcc tggctgactg agtttttcat agatgtactc 13380
tatcagggtt aggaagttcc cttttattcc taggttgttg agtctatttt atattacttt 13440
tttagagaca gtcttgctct gtccctcagg ctgtagcaca gtggctcaat catagctcac 13500
tgcagccttg aactcctagg ttcaagagat ccgcctgcct cagccttctt agtagctggg 13560
attacatgca tgcaccacca tactgggcta atttttttaa attttttata gagacaggg 13620
cttattacta tgttgcccag actggcattg agtcttttta tcattaatga gcaactgaatt 13680
ttgtcaagtg cttttataat acctattgtg atgatcatag ggttttgttc tttagtctac 13740
cgatacgcta tattgcatta agtgattttt ttgaatgtta aaccaacctt gcattttttt 13800
gggtgtataag tcttatttga tcaatgtgta ttatcctttt atatgggtgt ggatttagtt 13860
tgctactatt ttgttgagga tttttgtgtc tatattcata agagatatgt gtctgtagtt 13920
tcttgatgat tctttgtctg gttttagaat cagggtaatg ctggcctcat agaataaatt 13980
gggaagtgtt gtcttttcta tgtgatggga gagtttgtga atcattggta ttaatttttc 14040
tgtaaatgtt tggtagaatt cacaaataaa ggcactctgag cctgggctct tctttgtggg 14100
aagtttttgg cttttttttt ctttaaaaat tttcattgtg gctgggcatg gtggctcacg 14160
cctgtaatcc cagcactttg gggggccaag acgggtgaat cacctgaggn nnnnnnnnnn 14220

```



```

nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn 14280
nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn 14340
nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn 14400
nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn 14460
nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnngag tgcaatgatt gcagtcttgg 14520
ctcactgcaa cctctgccac ctgggctcaa gagattctcc tgcctcagcc tcctgagtag 14580
ctgggattac aggcattgccc caccatgccc agttaatttt tgtattttta gtagagacgg 14640
ggattctcca tgttgaccag gctgggtctct aactcctcac ctcaagtgat ccgcccgcct 14700
cggcctccca aagtgctggg attacaggca tgagccacca cgcccggcct taaaaatttt 14760
tttaattgtac agttgagtag tatttaatac attcacattg ttgtgtaccc agtttccaga 14820
actcttcate ctacagaact gaaactccat acccattaaa tgagtcccca ttctctttcc 14880
cccagctcat ggcaaacagc attctatttt cagtctctat gaatttgatt agtttagata 14940
cttcatactg taagtggaaat catatgggat ttgtctttta gtgactgcct aatttaaaaa 15000
aaattttttt gagacggagt cctgctctgt cgcccaggct ggagtgcagt ggcaccatct 15060
ctgctcactg caacctccac ctcccagggt caagtgattg tcctgcctcg gcctcccacg 15120
tagctgggat tacagggtgt cgccacaaca cccggctaatt ttttgtattt ttaggtagag 15180
actgggtttc accatgttgg ccaggctggt ctogaactcc tgacctcaaa ttatccacct 15240
gccttggcct cccaaagtgc tgggattaca ggcgtgagcc actgtgcca gcctccatgt 15300
tgttttttcac aacacctgta tcattttacat ttccaccaac agtacacaag aatttcagtt 15360
tctccacatc cttgctagca gttgttatta tctgtttttt tttaattggt tcttttttcc 15420
tttttctttt tttttttttt tgagacgggtc ttattcttgc tcatgctgga gtgcagtggg 15480
gcaatgtgat agctcactgc agcctcaacc tctgggctca agcagtcctg ccacctcagc 15540
ctccacatag gtgggactgc aggtgtgcac caccacttgt ggctaattta aaaaattttt 15600
tcgtagagac agagtctcac agtggtaccc aggtggtct tgaacttctg agcacaagtg 15660
atcctccac ctcagcctcc caaaataatg agattagaga catgagccaa catgcccac 15720
cagttttggt tgtttggttt gttttgtgtt tttgagacag agtctcactc tattgcccag 15780
gctggagtgc aacggcatga tcttagctca ctgcaacctc cgcctcccag gttcaagtga 15840
ttctcatgcc ccagcctcct gagaagctgg gattacaggt gtaccaccac acccagttat 15900
ttttgtattt ttagtagaca tgggggtttg ccatgttggc caggctggtc ccgaactcct 15960
gacctcaagt gatctgctcc cctcagcctc ccaagggtgt agaattaagt ttttctttct 16020
ttctttcttt cttttttttt ttttttttga gacagagtct cactctgtca ccagggcagg 16080
agtgcattgg cacgggtctg gctcattgta acctctgcct ccagattca agtagtgatt 16140
ctcctgtctc agcctcccaa gtagctggga ttacaggcat gcaccaccac gccagctaa 16200
ttttttgtat ttttagtaga aacgggggtt caccatgttg gtcaggctga tctcaaactc 16260
ctgaccccaa gtgatccacc cgccttgccc tcccaaagtg ttgggattac aggcgtgagc 16320
cactgtgcct ggtttttattt ttattattat tatttttaat agttcctatt ctaatgggta 16380
tgaggtagtg aggtgggtgg ttgtgggtgt tttatgaatg tttaattgga aatgggtggc 16440
cattgtgtgc aggaaaaacc tcctaaattg tgtcaaactc ctggaaaatg aaatatcatt 16500
ccagttgcaa gaatatcttt tttttttttt ttttttttta agacagagtc tcaactctgtc 16560
accaggcggg agtgcagtgg cacgatctcg gctcactgca acctccgcct cctgggtccgc 16620
ctcccgggtt caagtgatcc ttctgcctta gcctcccaag tagctgggac tacaggcgcg 16680
tgccaccact cctggctaatt ttttttgtat ttttagtaga gatgggggtt caccatgttg 16740
gccaggatgg tctcaatctc ttgacctaga gatccgcctg cctcggcctc ccaaagtggg 16800
gggattacag gggggtcacc gtgcccagcc acaagaagat cttgagcatg tgaatgatca 16860
gaaatgattt agcctatgta ggcactaggc caggtagtga aattcaggga aaataattca 16920
gatgcttctg agctatcact tatgaactaa gaaacagctt aaagccatta tagtgtgttt 16980
cctgaagatg aaagcatatg gtaagatgaa atagtgatta ttttttaaaa attactactc 17040
cagaaaggaa aagtttacta atttttatta ctaaagttta ctgttggtgg gtgcggtggc 17100
tcacacctgt aatcccagca ctttgggagg ccgaggcagg cggtcacct gaggtcagga 17160
gttcgagacc agcctgacca atatggtgaa acccatctc tgctaaaaat aaaaaattag 17220
gccgggcgcg gtggctcatg cctgtaatcc cagcactttg ggaggccgag gcaggtggat 17280
cacgaggtca ggagattgag accatcctgg ctaacacggt gaaaccccgct ctctgctaaa 17340
aatacaaaaa tcagccgggc gtcttggcag gcacctgtag tcccaggtag tcaggagttt 17400
tgagacggga gaatggcgtg aaccgggaag gcggagcttg aagtgagccg agattgcgcc 17460
gcttcagtcc agcctggacg acagagtgag actctgtctc taaaaaaata aataaataaa 17520
aataaaaaat tagctgggtg tgggtggcag cacctgtact ccagctact cgggaggctg 17580
aggcaggaga attgcttgaa cccgggagat ggaggttgcg gggagccaag attgcgccac 17640

```

tgcactccag	cctggcgaca	gagtgagact	ctttctcaga	aaaaaatatg	ataattaaaa	17700
gttgagacgt	tcttcgccga	gagtggtcgg	ggtttcctgc	ttcaacagtg	cttgggacgga	17760
acccggcgct	cgctctgcac	cccggccggc	cgcccatagc	cagccctccg	tcacctcttc	17820
accatgccct	cggactgccc	caaggccccc	gccgcagctc	cagcgccgcg	tagccaccac	17880
tgccgctgcc	gccgcctctc	cttagtcgcc	ggcatgacga	ccgcgtctac	ctcgcagggtg	17940
cgccagaact	accaccaaga	ctcagaggcc	gccatcaacc	gccagatcaa	cctggagctc	18000
tacgcctcct	ccatttacct	gtgcgtggct	tactactttg	acagcgatga	tgtggctttg	18060
aagaactttg	ccaaataactt	tcttcaccaa	tctcatgagg	agaggggaaca	tgctgagaaa	18120
ttgatgaagc	tgtagaacca	acgagggtggc	cgaatcttcc	ttcaggatat	caagaaacca	18180
gactgtgcgg	ggagaatgcg	atgggagagc	gggctgaatg	cgatggatta	catttgga	18240
aaattgtgca	ttttgcatta	catttgga	aaaatgtgaa	tcagtcacta	ctggaactgc	18300
acaaactggc	cactgacaaa	aatgaccccc	atttgtgtga	cttcattgag	acacattacc	18360
tgaatgaaca	agtgaaggcc	atcaaagaat	tgggtgacca	cgtgaccaac	atgcacgaga	18420
tgggagcgcc	cgaatctggc	gtggcagaat	acctctttga	caagcacacc	ctgggagaca	18480
gtgataatga	aagctaagcc	tcaggcta	ttccccatag	ccatagggtg	acttaccttg	18540
tcaccaaggc	agcgcatgta	tgttggggtt	tcctttacct	tttctataag	ttgttccaag	18600
acaccactt	aagttctttg	atttgtacca	ttccttcaaa	taaataaatt	tggtaccctc	18660
cccccccca	aaaaaaaaat	gtactgtggg	ctggcgtagt	ggctcatgcc	taaatctcag	18720
cactttggga	ggctgaggcg	ggaggatcac	ctgagggtcgg	gagtttgaga	ccagcctggg	18780
caacatggtg	aaaccccggtg	tctactaaaa	atataaaaac	tagccagtca	tggtggcaca	18840
cacctgtaat	cccggctact	tgggaggctg	aggcatgaga	atcacttgaa	cctgggctgc	18900
ggaggttgta	gtaagctgag	atcatgccac	tgtactccag	cctgggtgac	agggagacat	18960
tctctctctc	aaaaaaaaaa	aaaaaaca	aaaaacaaaa	caaaccaaca	aaacaaagta	19020
atccaggaac	aacaacatga	tgaaggactg	catgcaggac	tcagtgatgg	atggtggaag	19080
acagccagga	agttaagcat	gactctggta	ttaagtgttg	tctgggagag	ttaagattcc	19140
atttacagaa	ataagacctg	taggggaagc	tcttgatttt	tttttttttg	cagactgctg	19200
atttcctgat	tacatgtgtt	aagtttgagg	tatagagaga	aagaacatcc	tggccgggtg	19260
cagtggctca	cacccgtaat	cccagcactt	tgggaggcca	agggtggcag	atcacgaggt	19320
ccaggagatc	gagaccatcc	tggccaacat	ggtgaaaccc	cgtctctact	aaaaatacaa	19380
aaattagctg	ggcgtggcgg	cgcgtgcctg	ttatcccagc	tactcaggag	gctgaggcag	19440
gagaattgct	tgaacccgag	aggcagaggt	tgtgatgagc	cgagatcgcg	ccactgcact	19500
ccaccctggc	aacagagcta	gactctgtct	caaaaaaaaa	aaaaagaaag	aaaaaaagaa	19560
catcctgtag	aaacaggcag	tcagagggtat	agaactacac	agaatccaag	agatctttca	19620
agaaaagtga	catgcagcaa	gagaaactat	caaggggggt	aacaacctat	agaatgggag	19680
aaaatattca	caaagtatac	atccaacaaa	agtctaata	ccaaaatcta	taaggaacct	19740
aacaagcaaa	aagcaaataa	cccccttaaa	aagtgggcaa	aggacatgaa	cagatacttc	19800
tcaaaagacg	tacatgtggc	ccacaaacat	gaaaaaacgc	ccatttctaa	tcacagaca	19860
aatgcaaatt	agaaccacaa	agagatacca	tctcacacca	gtcagaacag	cttttggttaa	19920
aatgtcaaaa	aatgagaaac	gttgggtgagg	ctgcagagga	aagcagacac	ttgtacactg	19980
ttggtgaagg	tgtaaattaa	tttagcgtag	gcacagtcag	tttggagatt	tctcagagaa	20040
ctaagagtgg	aactaccatt	agaccagca	atctgatggc	tggatatacg	gccaaaggaa	20100
aataaatcat	tctgcaaaaa	gaacatatgt	acctgtatgt	tcattgtggc	accattcaca	20160
atagcaaaaa	cattgaatca	actcatgtgc	ccatcagtgg	cggactagaa	aagaaaagaa	20220
aatatggtac	atagccatca	tgggaatacta	tgcacccatt	aaaaataatg	aaataatgtc	20280
tttgcaacaa	catgaatgta	gctggaggggc	attatcctaa	gcaaactaac	acagaaacag	20340
aaaaccaa	actgcgtgtt	ctcacgcagt	gagagtggga	gctgaacatc	aagtacacat	20400
ggatgtaaag	atggcaacaa	tagacatggg	tctactagag	gtggtggtgc	ggcagggtgg	20460
gggtgggggt	tgtgtggcag	aggaacagct	gaaaaactac	ctatttgata	ctatacccag	20520
cacctgggaa	acgggttcag	tcatacccca	aacctcagca	tcacacagta	tacctttcta	20580
acaaacttac	acatgtattc	tgtgattcta	aaataaacat	tgaaaaataa	aaaaaaaact	20640
gacatgggtt	gtactgttta	atctgacata	atggctaggg	gaaatgaagt	ctgcagaatg	20700
gctgttttacg	gatgttggtg	ttgttggtga	gatgagggtc	cactatgttg	cccaggcttg	20760
aactcctggc	ctcaagcagt	cctcctgcct	tgacctccca	aaatgttgag	attacatgca	20820
tgagccattg	ccaaaacggc	tatttggtatt	gctgttaagg	ttattacatt	ctctgtgtag	20880
taagaccttg	aaggagaagg	atttgagatc	aggagtttaa	gaaaaaatgt	taatctagga	20940
agagaggata	atttctgtgt	ttggccagtt	gcaatggctc	acgcctgtaa	tcccagcgtt	21000
ttgggaggcc	gagctgggca	gatcacttga	gctcaggagt	ttgagaagag	cctgggcaac	21060

atggtgaagc cccgtctcta ctgaaaatac aaaaattggc tgggcgtggt ggcaggtgcc 21120
cactgtaatt tcagccactc aggagactga ggcaggagaa ttgcttgaac ctggaaagtg 21180
gaggatatgg tgagcctaga tcgcgccact ctactccagc ctgggagact ccatctcaaa 21240
aaaaaaagaa aacaagatgc tgaaatgaag taattaccac agtcaatgtg atcctataac 21300
tttgttttct tttagagatg gggctctccct ctgtcaccca ggctggagtg cagtgggtgca 21360
tcatagcttt ctgcagcctc cacctcctgg gctcaggtgg tcctcttgcc tcagtgttcc 21420
gagtagttag gactgactgc aggtgcatgc tgctatgcct ggctaacttt aaaatttttt 21480
tgtagaggcg gggctcttgc atgttgccta ggctgggtct caactcctga tctcaatcag 21540
tcctcctgcc taccttccca ggcgctggg attacaggtg tgagccatcg cacctagcca 21600
atctcataac attttatgac tagcaaactt agtagttctg attcagggcat aaatcagttg 21660
gtgggggttat acaagggttg gtgagttttt ctagatttct aagagaccat gttgaaatac 21720
ttggctctgt ctgagtaagg gatagagaga agcaaagggtg tgggtaaagg ttatgagcag 21780
acatgtaaag ggggcaaatt aaagtgttta ggggaagggtg aacagttcca aatcataata 21840
tagatcccaa gcctcaccaa gaagtgaagg agagggtaat tgtgcgatag ttctcaggac 21900
tgagacctca aggtgtaaga cgaatcttta ttgtgggtgg tcctctttga gaaaaagaac 21960
aaaaaagaaa aatgtgaaat ggggtgctaaa tttcagagca gaatacat atatgtgtat 22020
gtgtgtgtgt atatatatat atatttttat gtagcaaatt ccagtttaca aagggttttt 22080
atgtgttttg catcattaca acagttctgt gatgatgtgg atgtgggtgga tattgtaatt 22140
ccacatccca gaggataaaa ttgaggcaca ccataaggta gctgaccaga gatcatgcag 22200
tatatggtag aatggcaact tgaggccagt tcttaggtgt gtttggtgtc cttattcaga 22260
atggaacaat gtggtttatt gtacaaaaat ttaaaaatga atgtcgaaaa gtagaaatta 22320
ttacccaaat ctcacctct gtggttgttt gctatgtgat cttccagact cacatataca 22380
ttagataaat tttttttttt ttttttgaga cggagtcttg ctctgtcgcc ctggcnnnnn 22440
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22500
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22560
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22620
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22680
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22740
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22800
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22860
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22920
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 22980
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23040
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23100
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23160
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23220
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23280
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23340
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23400
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23460
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23520
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23580
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23640
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23700
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23760
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23820
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23880
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 23940
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24000
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24060
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24120
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24180
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24240
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24300
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24360
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24420
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 24480

nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24540
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24600
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24660
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24720
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24780
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24840
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24900
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	24960
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnncaaaaat	tagctgggag	25020
tagtggtggg	cgctgtggt	cccagctact	tgggcagctg	aggcaggaga	atcgcttgaa	25080
cctgggaggt	ggaggttgca	gtgagccaag	accatgccac	tgcactccag	cctgggcaac	25140
agagcaagac	tcttgtctta	aaaaacaaaa	aagtgtaccc	agttgagctg	attctttatc	25200
tttttttcac	tggagaacta	agtatacagg	tgagaaaaga	cgagatattt	ataccgaga	25260
gaattgatgg	tgaatccat	ttttttggat	cagaacttcc	ccaaacagtg	tccttcaa	25320
agggttcagg	ggtgctaaga	tatttatccc	ctcaaccctt	ggggttcc	ccagtatggc	25380
atataaatat	tgtatcactt	tctatgtgtg	gggagcagtg	ctccaggtga	ccttccttcc	25440
tttcttctta	ggggaggaag	ttgccttgta	ctgtgccaaa	tatcttctg	atatcatcaa	25500
agatcagaag	gcctacaagg	aaggcaagct	acagaaggct	tgtctgctta	caccgccc	25560
tcctcacttg	tgtaggcttt	tcccttggtc	tctagccctt	gggcttttcc	tttctttttg	25620
tcctctagct	gctgctgctt	atttactctt	gaagaattct	gttccataaa	cgagcttatt	25680
ggccgccttt	tagacttgcc	ttattattcc	taggcctctg	agctgttttt	atctgtgagt	25740
gtctcttagt	gtggtggctc	acactcttaa	tttgtattcc	atccttggtg	tcaggattgt	25800
atataggagg	ttcattttgt	actagtctta	gactattttg	cttataattc	ggcttttaga	25860
gatgccttct	tggctattga	cgccaaattg	accactgaag	aagtcattaa	agagctggca	25920
cagattgcag	ggcgaccac	tgaggatgaa	gatgaaaaag	aaaaagtagc	tgatgaagat	25980
gatggtgagt	gtggcatccc	ttgtttgagg	ggaaatcagc	attttaagaa	atattcttta	26040
atattactta	tcaattctaa	gataggatgg	ctttctaggg	acctggggag	tccttatggt	26100
aaagaaacct	atgatgttct	cctgcattgt	atgtgggttat	gaaaaggagg	gagagaatta	26160
tcttcgttga	gtggcatctg	agctgtaagc	attgtatata	cattatcttt	tgtcattgtg	26220
atggggtctt	cctgggttct	gctagtattt	atgtgctttt	ttttccctc	aagactggag	26280
cagttattag	ccccaatagc	caatcattaa	gcctaaatcc	taattcacag	tagcattgtg	26340
ggcttcctgg	atcctcagcc	agaatagggt	ttttacaact	taacaataaa	aaatgagacg	26400
tcagagggga	agtatagtaa	ctagtgttgt	tttgattaag	aaggggatga	aacacaaaaa	26460
ccaaaagaag	tctgtggagg	aggaggagct	agggcattgt	cttctgagac	ttgagcgaga	26520
ggaaccttgg	gagtgggagg	ttgtggggaa	gttagaggct	gcaagggtg	ttgaggtagt	26580
gagagggacg	gatcccatga	ggagtctggc	atgggggtct	tgatttagcc	tcttccttgc	26640
agtggacaat	gaggaggctg	cactgctgca	tgaagaggct	accatgacta	ttgaagagct	26700
gctgacacgc	tacgggcaga	actgtcacia	gggcccctcc	cacagcaaat	ctggagggtg	26760
gacaggcgag	gaaccagggt	cccagggcct	caatggggag	gcaggacctg	aggactcaac	26820
tagggaaact	ccttcacaag	aaaatggccc	cacagccaag	gcctacacag	gcttttcctc	26880
caactcggaa	cgtgggactg	aggcaggcca	agttggtgag	cctggcattc	ccactggtga	26940
ggctgggcct	tcctgctctt	cagcctctga	caagctgcct	cgagttgcta	agtccaagtt	27000
ctttgaggac	agtgaggatg	agtcagatga	ggcgaggagg	gaagagggaag	acagtgaggt	27060
aagggcctgt	gagggcaggc	agatgctgaa	gttgacagaga	ggctctgttt	ggttgccgtc	27120
tgtagttttc	aactctcttt	ccttctccta	ttttgacatc	atcccccaag	accactgta	27180
ttctaagctt	tagtcttgaa	ttcattgagc	tccatcatca	caggtaccat	ttgccttttt	27240
acctcttctt	ttgttggtac	tataacaagc	agatctagtt	ctggcttttc	agagtctgtc	27300
tcctagagag	agaacaagga	gatagttgtt	accttggtga	gttgactgtt	ttcttctctg	27360
gaaaatttat	tttctggcca	cagtgcctga	aagatatattt	tggctggcag	cccttgccct	27420
gtcctgggct	ttttgctagt	gactgctaag	cccagttcag	gatgtcagtt	gtactcatgc	27480
tagccctttc	catcccccca	attttcatga	ccatataact	gtatctttca	gtgttttgag	27540
gacctgtgtt	cagtcaggac	ctcttgattc	tgagtatgag	ctgtggggag	ggaggggatc	27600
atcccagctt	cagcagctctg	ggatcctccc	cctggcaggga	atgcagcgag	gaagaggatg	27660
gctacagcag	tgaggaggca	gagaatgagg	aagatgagga	tgacaccgag	gaggctgaag	27720
aggacgatga	agaagaagaa	gaagagatga	tgggtgccagg	gatggaaggc	aaagaggagg	27780
tgtgtgggga	aggggagcaa	tgagtcttga	aaagccacaa	ggcagggtgtg	aatcccctaa	27840
ttttgatattt	gagacagggg	atccccctga	tacttttagga	tggaaagta	agtcagggg	27900

```

atttattctg caaggggaat gagatggtaa gcctttgggg ttgaattatc taaaaacaag 27960
ggagagggag tgtgctgctg tctctagaaa gatgaaatgt gtgcttctcc tgtttggttaa 28020
agctcttttg ggggtcccag tgaaagcaag cataggtgaa cgatcaggag cacatcagtg 28080
aggaacgcat gttcagaagc ccccatgatg ctctttttct tcctcttaag cctggctctg 28140
acagtggtag aacagcggtg gtggccctga tacgagggaa gcagttgatt gtagccaacg 28200
caggagactc tcgctgtgtg gtatctgagg ctggcaaagc tttagacatg tcctatgatc 28260
acaaaccaga ggatgaagta gaactagcac gcatcaagaa tgctgggtggc aaggtcacca 28320
tggatgggag agtcaacggg ggcctcaacc tctccagagc cattggtaag ggccaagaaa 28380
ctgggaaaga nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 28440
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 28500
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 28560
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 28620
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 28680
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnatgggtgaa 28740
accccatctc tactaaaaaa aaaaaaaaat acaaaaatta gctgggcatg gtagtgcaag 28800
cctgtaatcc cagctactca ggaggctgag gcaggagaac cgcttgaatc cgggaggcgg 28860
aggttgtagt gagccgagat cgtgccatta cactccagcc tgggctacaa gagtgaact 28920
ccgtctcaaa aaaaaaaaaa caaaaaagac ttaaaataaa aagaccagtg agtgactttc 28980
ttaaggttca gcagtctggt ggcagggttg aaactagaaa aactaggact taggactcag 29040
ttccccattc cactagatta tggaaactttg taaagaaggg aaatgaatgg caaggtttga 29100
cctgccacaa acacaagtct gtgggaagta tccaaactgc tcatcaacca ttcctttact 29160
ccaggggacc acttctataa gagaaacaag aacctgccac ctgaggaaca gatgatttca 29220
gcccttctg acatcaaggt gctgactctc actgacgacc atgaattcat ggtcattgcc 29280
tgtgatggca tctgggtgagc actggcagaa tgccctaaat tccctttct gcagcatgtc 29340
ttctcttata ggactcaggg cacctctagg attagagcct aggcagacct aggcctcttg 29400
gtgggtgaag agcacccaga ctaaggcaga gctgagaatt tctgtagtta tttacactgg 29460
cctgggccac cacctctgtc catactcctc tacgctgcct tagtgagact ggaagattct 29520
gactgttggt cttgaccca ggaatgtgat gagcagccag gaagttgtag atttcattca 29580
atcaaagatc agccagcgtg atgaaaatgg ggagcttcgg ttattgtcat ccattgtgga 29640
agaggtgagt accaggggtg agaagagagg gtgtctggtc tgcacagcca ggggtt 29695

```

<210> 4
 <211> 146
 <212> PRT
 <213> Human

<400> 4
 Gly Asp His Phe Tyr Lys Arg Asn Lys Asn Leu Pro Pro Glu Glu Gln
 1 5 10 15
 Met Ile Ser Ala Leu Pro Asp Ile Lys Val Leu Thr Leu Thr Asp Asp
 20 25 30
 His Glu Phe Met Val Ile Ala Cys Asp Gly Ile Trp Asn Val Met Ser
 35 40 45
 Ser Gln Glu Val Val Asp Phe Ile Gln Ser Lys Ile Ser Gln Arg Asp
 50 55 60
 Glu Asn Gly Glu Leu Arg Leu Leu Ser Ser Ile Val Glu Glu Leu Leu
 65 70 75 80
 Asp Gln Cys Leu Ala Pro Asp Thr Ser Gly Asp Gly Thr Gly Cys Asp
 85 90 95
 Asn Met Thr Cys Ile Ile Ile Cys Phe Lys Pro Arg Asn Thr Ala Glu
 100 105 110
 Leu Gln Pro Glu Ser Gly Lys Arg Lys Leu Glu Glu Val Leu Ser Thr
 115 120 125
 Glu Gly Ala Glu Glu Asn Gly Asn Ser Asp Lys Lys Lys Lys Ala Lys
 130 135 140
 Arg Asp
 145

<210> 5
 <211> 139
 <212> PRT
 <213> Human

<400> 5
 Met Gly Ala Tyr Leu Ser Gln Pro Asn Thr Val Lys Cys Ser Gly Asp
 1 5 10 15
 Gly Val Gly Ala Pro Arg Leu Pro Leu Pro Tyr Gly Phe Ser Ala Met
 20 25 30
 Gln Gly Trp Arg Val Ser Met Glu Asp Ala His Asn Cys Ile Pro Glu
 35 40 45
 Leu Asp Ser Glu Thr Ala Met Phe Ser Val Tyr Asp Gly His Gly Gly
 50 55 60
 Glu Glu Val Ala Leu Tyr Cys Ala Lys Tyr Leu Pro Asp Ile Ile Lys
 65 70 75 80
 Asp Gln Lys Ala Tyr Lys Glu Gly Lys Leu Gln Lys Ala Leu Glu Asp
 85 90 95
 Ala Phe Leu Ala Ile Asp Ala Lys Leu Thr Thr Glu Glu Val Ile Lys
 100 105 110
 Glu Leu Ala Gln Ile Ala Gly Arg Pro Thr Glu Asp Glu Asp Glu Lys
 115 120 125
 Glu Lys Val Ala Asp Glu Asp Asp Val Asp Asn
 130 135

<210> 6
 <211> 145
 <212> PRT
 <213> Mus musculus

<400> 6
 Gly Asp His Phe Tyr Lys Arg Asn Lys Asn Leu Pro Pro Gln Glu Gln
 1 5 10 15
 Met Ile Ser Ala Leu Pro Asp Ile Lys Val Leu Thr Leu Thr Asp Asp
 20 25 30
 His Glu Phe Met Val Ile Ala Cys Asp Gly Ile Trp Asn Val Met Ser
 35 40 45
 Ser Gln Glu Val Val Asp Phe Ile Gln Ser Lys Ile Ser Gln Arg Asp
 50 55 60
 Glu Asn Gly Glu Leu Arg Leu Leu Ser Ser Ile Val Glu Glu Leu Leu
 65 70 75 80
 Asp Gln Cys Leu Ala Pro Asp Thr Ser Gly Asp Gly Thr Gly Cys Asp
 85 90 95
 Asn Met Thr Cys Ile Ile Ile Cys Phe Lys Pro Arg Asn Thr Val Glu
 100 105 110
 Leu Gln Ala Glu Ser Gly Lys Arg Lys Leu Glu Glu Ala Leu Ser Thr
 115 120 125
 Glu Gly Ala Glu Asp Thr Gly Asn Ser Asp Lys Lys Lys Ala Lys Arg
 130 135 140
 Asp
 145

<210> 7

<211> 139

<212> PRT

<213> Mus musculus

<400> 7

Met	Gly	Ala	Tyr	Leu	Ser	Gln	Pro	Asn	Thr	Val	Lys	Cys	Ser	Gly	Asp
1				5					10					15	
Gly	Val	Gly	Ala	Pro	Arg	Leu	Pro	Leu	Pro	Tyr	Gly	Phe	Ser	Ala	Met
			20					25					30		
Gln	Gly	Trp	Arg	Val	Ser	Met	Glu	Asp	Ala	His	Asn	Cys	Ile	Pro	Glu
		35					40					45			
Leu	Asp	Asn	Glu	Thr	Ala	Met	Phe	Ser	Val	Tyr	Asp	Gly	His	Gly	Gly
	50					55					60				
Glu	Glu	Val	Ala	Leu	Tyr	Cys	Ala	Lys	Tyr	Leu	Pro	Asp	Ile	Ile	Lys
65					70					75					80
Asp	Gln	Lys	Ala	Tyr	Lys	Glu	Gly	Lys	Leu	Gln	Lys	Ala	Leu	Gln	Asp
			85						90					95	
Ala	Phe	Leu	Ala	Ile	Asp	Ala	Lys	Leu	Thr	Thr	Glu	Glu	Val	Ile	Lys
			100					105						110	
Glu	Leu	Ala	Gln	Ile	Ala	Gly	Arg	Pro	Thr	Glu	Asp	Glu	Asp	Asp	Lys
		115					120					125			
Asp	Lys	Val	Ala	Asp	Glu	Asp	Asp	Val	Asp	Asn					
	130						135								